

AdS_3 ASYMPTOTIC (SUPER)SYMMETRIES

Karin Bautier

Physique Théorique et Mathématique

Université Libre de Bruxelles, Campus Plaine C.P. 231

Boulevard du Triomphe, B-1050 Bruxelles, Belgium

Abstract The Chern-Simons formulation of AdS_3 supergravity is considered. Asymptotic conditions on the Rarita-Schwinger fields are given. Together with the known boundary conditions on the bosonic fields, these ensure that the asymptotic algebra is the superconformal algebra, with the same central charge as the one of pure gravity. It is also indicated that the asymptotic dynamics is described by super-Liouville.

It has been pointed out in [1] that the asymptotic symmetry group of anti-de Sitter gravity in three dimensions is the conformal group in two dimensions with a central charge $c = 3l/2G$. This result was obtained by working out explicitly the boundary conditions and solving the asymptotic Killing equations [1]. It has been shown in [2] that the boundary dynamics at infinity is described by Liouville theory up to terms involving the zero modes and the holonomies that were not worked out. In the following, we will use the Chern-Simons formulation of AdS_3 (1,1)-supergravity and extend the analysis of [1, 2] to the supersymmetric case. This was originally presented in [3] written in collaboration with M. Bañados, O. Coussaert, M. Henneaux and M. Ortiz and we refer to it for more details.

Chern-Simons action. AdS_3 (1,1)-supergravity can be written as a Chern-Simons theory [4]. The relevant group is $OSp(1|2) \times OSp(1|2)$ and the action is:

$$I[A, \psi; \tilde{A}, \tilde{\psi}] = I[A, \psi] - I[\tilde{A}, \tilde{\psi}] \quad (1.1)$$

*ULB-TH-99/16

where $I[A, \psi]$ and $I[\tilde{A}, \tilde{\psi}]$ are the Chern-Simons actions for the supergroup $OSp(1|2)$:

$$I[A, \psi] = \frac{k}{4\pi} \int \left[\text{Tr}(AdA + \frac{2}{3}A^3) + i\bar{\psi} \wedge D\psi \right] \quad (1.2)$$

(with a similar expression for $I[\tilde{A}, \tilde{\psi}]$). The constant k is related to the 3D Newton constant G and the anti-de Sitter radius l through $k = l/4G$. The action (1.2) is invariant under the bosonic gauge transformations $\delta_\lambda A = D\lambda$, $\delta_\lambda \psi = -\lambda\psi$ and under the fermionic ones $\delta_\rho A^a = i\bar{\rho}\gamma^a\psi$, $\delta_\rho \psi = D\rho$, where the gauge parameters $(\lambda^a, \rho) \in osp(1|2)$.

Boundary conditions. The boundary conditions at infinity on the bosonic fields for an asymptotically AdS space have been given in [1] in the metric representation and they were reexpressed in the connection representation in [2]. To supplement these conditions by appropriate boundary conditions on the fermionic fields, one follows the procedure of [5]: one starts with the known physical metrics that should be included in the theory - here, the black hole solutions [6] - and acts on them with the anti-de Sitter supergroup. One then obtains, for one $OSp(1|2)$ copy, the following boundary conditions:

$$A_v = 0, \quad \psi_v = 0, \quad (1.3)$$

$$A_r = b^{-1}\partial_r b, \quad \psi_r = 0 \quad (1.4)$$

and

$$lA_u = b^{-1} \begin{pmatrix} 0 & L/k \\ 1 & 0 \end{pmatrix} b, \quad l\psi_u = b^{-1} \begin{pmatrix} Q/k \\ 0 \end{pmatrix} \quad (1.5)$$

asymptotically. Here, $u = t + l\phi$, $v = t - l\phi$ and $L = L(t, \varphi)$ and $Q = Q(t, \varphi)$ are arbitrary functions which will be shown to be equal to the generators of the super-Virasoro algebra. The group element $b(r)$ is equal to

$$b(r) = \begin{pmatrix} (r/l)^{1/2} & 0 \\ 0 & (r/l)^{-1/2} \end{pmatrix} \quad (1.6)$$

The fact that ψ_u has one zero component implies that it is an eigenstate of the radial γ -matrix, which makes the induced spinors chiral in two dimensions. The other $OSp(1|2)$ fields satisfy analogous boundary conditions, with u and v interchanged, and depends on two additional functions $\tilde{L}$ and $\tilde{Q}$. For positive values of L_0 and $\tilde{L}_0$, the boundary conditions (1.5) represent a black hole, with $M = (2/k)(L_0 + \tilde{L}_0)$ and $J = (2l/k)(\tilde{L}_0 - L_0)$. Anti-de Sitter space corresponds to $L/k = \tilde{L}/k = -1/4$.

Asymptotic symmetry. The most general supergauge transformations that preserve the boundary conditions (1.3), (1.4) and (1.5) are characterized by gauge parameters (λ^a, ρ) that must fulfill, to leading order,

$$\lambda(u, r) = b^{-1}\eta(u)b, \quad \rho(r, u) = b^{-1}\varepsilon(u) \quad (1.7)$$

with

$$\eta^+ = \frac{\eta^- L}{k} - (1/2)(\eta^-)'' + \frac{iQ\epsilon}{2k}, \quad (1.8)$$

$$\eta^1 = -(\eta^-)', \quad (1.9)$$

$$\varepsilon = \begin{pmatrix} -\epsilon' + \eta^- Q/k \\ \epsilon \end{pmatrix}, \quad (1.10)$$

where $'$ denotes derivative with respect to the argument, and η^- and ϵ are two residual functions of u . The asymptotic symmetry acts on the components L and Q of the connection that remain at infinity in the following way:

$$\delta L = (\eta^- L)' + (\eta^-)'L - \frac{k}{2}(\eta^-)''' + \left(\frac{iQ\epsilon}{2}\right)' + iQ\epsilon', \quad (1.11)$$

$$\delta Q = -k\epsilon'' + L\epsilon + (\eta^- Q)' + \frac{1}{2}(\eta^-)'Q. \quad (1.12)$$

This indicates that L and Q form a super-Virasoro algebra, with a central charge c equal to $6k$ ($c/12 = k/2$).

To express this asymptotic symmetry in terms of Poisson brackets, we note that the canonical generators of the gauge transformations are:

$$G(\lambda^a) = \int_{\Sigma} \lambda^a \mathcal{G}_a + B, \quad S(\rho) = \int_{\Sigma} \bar{\rho} \mathcal{S} + F. \quad (1.13)$$

The boundary terms B and F must be chosen so that the generators G and S have well-defined functional derivatives [7]. Taking into account the above asymptotic conditions and gauge parameters, one obtains:

$$B = \frac{1}{2\pi} \int_{\partial\Sigma} \eta^- L, \quad F = \frac{-i}{2\pi} \int_{\partial\Sigma} \epsilon Q. \quad (1.14)$$

In the reduced phase space (where the constraints are zero), the generators of the asymptotic symmetry reduce to their boundary term (1.14). From equations (1.11) and (1.12), we compute their Poisson brackets and we get, after Fourier transformation, the asymptotic super-Virasoro algebra:

$$[L_m, L_n] = (n - m)L_{n+m} + \frac{k}{2}n^3\delta_{n+m,0} \quad (1.15)$$

$$[L_m, Q_n] = \left(\frac{1}{2}m - n\right) Q_{m+n} \quad (1.16)$$

$$\{Q_m, Q_n\} = 2L_{m+n} + 2km^2\delta_{m+n,0} \quad (1.17)$$

with a central charge equal to $c = 6k = 3l/2G$.

Dynamics at infinity. Following [2] and referring to [3] for details and references, we note that the Chern-Simons theory under the boundary conditions (1.3) induces the chiral Wess-Zumino-Witten model at the boundary. The other boundary conditions (1.5) turn out to be precisely the constraints that reduce the WZW theory based on the supergroup $OSp(1|2)$ to chiral $2D$ supergravity. The two chiral theories are combined to get the non chiral super-Liouville theory (up to zero modes and holonomies). This has been checked at the level of the action, using the Gauss decomposition for $OSp(1|2)$.

Acknowledgments

I would like to thank my collaborators M. Bañados, O. Coussaert, M. Henneaux and M. Ortiz, and the organizers of the Cargèse 99 ASI for a very pleasant school and for financial support. This work has been partly supported by the “Actions de Recherche Concertées” of the “Direction de la Recherche Scientifique - Communauté Française de Belgique” and by IISN - Belgium (convention 4.4505.86). The author is “Chercheur F.R.I.A.” (Belgium).

References

- [1] Brown, J.D. and Henneaux, M. (1986) *Commun. Math. Phys.*, **104**, pp. 207-226.
- [2] Coussaert, O., Henneaux, M. and van Driel, P. (1995) *Class. Quant. Grav.*, **12**, pp. 2961-2966, gr-qc/9506019.
- [3] Bañados, M., Bautier, K., Coussaert, O., Henneaux, M. and Ortiz, M. (1995) *Phys. Rev.*, **D58** 085020, hep-th/9805165.
- [4] Achúcarro, A. and Townsend, P.K. (1986) *Phys. Lett.*, **B180**, p. 89.
- [5] Henneaux, M. and Teitelboim, C. (1985) *Commun. Math. Phys.*, **98**, pp. 391-424.
- [6] Bañados, M., Teitelboim, C. and Zanelli, J. (1992) *Phys. Rev. Lett.*, **69**, pp. 1849-1851, hep-th/9204099.
- [7] Regge, T. and Teitelboim, C. (1974) *Ann. Phys.*, **88**, p. 286.